

Emergency Neurological Life Support Spinal Cord Compression Protocol Version 5.0

Authors

Safdar A. Ansari, MD
Jennifer M. MacDonald, MD, PhD
Stephanie Qualls, RN, BSN, CNRN
Ryan Martin, MD

Last updated: May 2022

Spinal Cord Compression Algorithm




Table of Contents

Spinal Cord Compression Algorithm	3
Checklist.....	5
Communication	6
Acute Disc Herniation.....	7
Decompression.....	7
Acute Hemorrhage.....	8
Reverse coagulopathy and decompression	8
Assess Airway and Hemodynamics	9
Cervical myelopathy may affect diaphragm.....	9
Corticosteroids	10
Empirical treatment if cancer is suspected	10
Emergent Transfer	11
To a facility that has spine imaging available	11
To a facility that has pediatric specialists	11
Empirical Antibiotics	11
Empirical treatment for presumed infectious cause.....	11
Imaging Not Available	12
No MRI or CT.....	12
Infectious Lesion	13
Empirical antibiotics and ESR	13
MRI Spine.....	14
Spine imaging is available	14
Negative Imaging or Intrinsic Spinal Cord Lesion	15
ENLS Non-traumatic Weakness Protocol.....	15
Neoplastic Disease	16
Steroids, decompression, radiation oncology.....	16
Paraplegia	17
Quadriplegia.....	18
Special airway issues	18
Spinal Cord Compression	19
Suspected myelopathy	19
Suspicion of Cancer	20
Possible metastasis.....	20
Suspicion of Infection	21
Consider epidural abscess	21

Clinical Checklist

- Brief history of the patient
- Spinal motion restriction (motion restriction) and pressure point padding
- Ensure proper ventilation, especially in the presence of quadriplegia
- Labs: CBC, Chemistries, INR/PT, PTT, platelet function assay (e.g., VerifyNow platelet reactivity profile)
- Obtain emergent spine imaging (MRI unless contraindicated)
- Alert spine surgeon
- Suspected neoplastic disease: administer corticosteroids (dexamethasone 10 mg IV load and 4 mg every 6 hours IV/PO), contact radiation oncology
- Suspected epidural infection: check ESR and start antibiotics
- Pediatric consideration: Dexamethasone dosing 1-2mg/kg to max 10mg bolus and 0.25-0.5mg/kg every 6 hours IV/PO. Ensure adequate perfusion (mean arterial pressure > 50th percentile for age = 55 + [1.5 x age in years])
- Initiate interfacility transfer if anything cannot be performed at your facility

Communication Checklist

- Age, gender, premorbid conditions and risk factors
- Onset and duration of symptoms
- Paraplegia or quadriplegia
- Spinal level of clinical involvement
- Vital signs
- Airway status
- Bowel or bladder involvement
- Results of lab tests
- Results of imaging studies, if available
- Medical history (i.e. cancer, intravenous drug abuse, immunosuppression)
- Therapies initiated
- Discuss further interventions to start now

Sample Sign-off Narrative:

“I am transferring to you a patient who I suspect has a spinal cord compression secondary to an epidural abscess.

He is a 67-year-old male with a history of hypertension, diabetes, and intermittent intravenous drug use who presented one hour ago with mid-back pain over the last two weeks.

Over the last four days, his legs have been weak, and today he was unable to walk.

He appears paraplegic with a T-10 sensory level.

He has had low-grade fevers and chills since his back pain started.

On exam, he is afebrile, blood pressure is within normal limits and he has no evidence of respiratory insufficiency. Strength in his arms is normal, but he is unable to lift his legs off the bed. His bladder was distended and a Foley was placed.

CBC and Coags are normal. ESR is pending. We do not have any spinal imaging available.

Given his history of IV drug use and recent fevers and chills, we suspect this is an epidural abscess leading to spinal cord compression.

Empirical antibiotics were started, including vancomycin, ceftriaxone and metronidazole.

We are initiating transfer to your facility for emergent MRI of the spine and surgical consultation.

Is there anything else we should do before transfer?”

Degenerative Spine Disease / Acute Disc Herniation

Decompression

Degenerative Spine disease with or without disc herniation is the most common cause of Spinal Cord Compression and presents in the cervical and thoracic spine due to flexibility of these segments. Back pain is typically worse when standing/sitting and improved in the recumbent position. Spine MRI imaging is the diagnostic modality of choice. Surgical decompression has been shown to improve patient outcomes and corticosteroids have uncertain therapeutic benefit.

If imaging reveals disc herniation or bone/osteophyte compression (spinal stenosis) of the spinal cord or cauda equina; this represents a neurosurgical emergency.

Acute Hemorrhage

Reverse coagulopathy and decompression

Spinal hematomas can present with rapidly developing paraparesis or quadriparesis with local or radicular pain. Spinal epidural hematomas are the more common compressive etiology, but intra-medullary hemorrhage (hematomyelia) can also occur and presents similarly. The most common causes of non-traumatic spinal cord hematoma include coagulopathy, vascular malformations, inflammatory myelitis, spinal tumors, and syringomyelia. Traumatic causes include penetrating injuries and high-impact, blunt-force trauma, iatrogenic post-surgical bleeding or a late complication after radiotherapy.

MRI with and without gadolinium is the modality of choice for diagnosing spinal hematoma, as it will demonstrate both the hematoma and any associated underlying pathology, including spinal cord edema.

Treatment includes rapid reversal of coagulopathy and urgent evaluation for surgical decompression.

Assess Airway and Hemodynamics

Cervical myelopathy may affect diaphragm

Patients with quadriplegia should be monitored closely for respiratory distress and failure.

- Frequent bedside pulmonary function testing and the detection of dysphonia and tachypnea are vital to diagnose impending respiratory collapse.
- Consider intubation if:
 - Forced Vital Capacity (FVC) < 1L or 20 mL/kg
 - Negative Inspiratory Force (NIF) < -30 cmH₂O
- The patient's own assessment of their respiratory status is frequently accurate but may be unreliable in the presence of analgesia or sedation.
- A low threshold should be maintained for placement of a definitive airway and mechanical ventilation, particularly if rapid progression of motor weakness is observed.

Spinal cord compression can cause hypotension (neurogenic) which may exacerbate spinal cord perfusion and induce secondary injury. Hemodynamic assessment should lead to urgent management of hypotension with normal blood pressure goals.

If there is any suspicion of trauma, do not extend the spine for intubation, and refer to the **ENLS protocol for Traumatic Spine Injury**.

For patients with total body weakness and cranial nerve deficits, consider a generalized neuromuscular disorder, or a stroke. Secure the airway first, then pursue the **ENLS protocol for Acute Non-Traumatic Weakness**, or the **ENLS protocol for Acute Ischemic Stroke**.

Corticosteroids

Empirical treatment if cancer is suspected

Patients with malignant SCC and acute neurologic dysfunction should receive corticosteroids. Empirical treatment with corticosteroids is recommended in patients with known malignancy and acute SCC, even if unconfirmed by MRI spine imaging.

- Steroids are given to reduce edema and decrease the chance of spinal cord venous infarction. The use of steroids in patients with compression from epidural metastatic disease is considered to be part of standard medical therapy.
- Given the safety profile and efficacy of the lower dose of dexamethasone, we recommend a dexamethasone 10 mg IV loading dose followed by 4 mg oral/IV every 6 hours as maintenance.

Emergent Transfer

To a facility that has spine imaging available

Due to the rapid progression of acute SCC syndromes, transfer agreements between emergency departments and acute care facilities should be pre-established to avoid prolonged attempts to find a facility when time is of the essence. Emergent transfer is warranted when the facility treating the SCC patient is unable to provide definitive care.

If emergent transfer is not feasible, or will be delayed significantly, then care should focus on spinal motion restriction, empirical treatment with corticosteroids or antibiotics (for the most likely diagnosis) and supportive care measures. The medical complications of spinal cord injury will need to be monitored and treated, including respiratory and hemodynamic instability, constipation, urinary retention, pain, deep vein thrombosis, and pressure ulcers.

To a facility that has pediatric specialists

Due to the unique etiologies of pediatric spinal cord compression and their treatments, transfer to an acute care center with pediatric specialists available should be arranged as quickly as possible. To prevent further injury, the spine should be immobilized and adequate ventilation and perfusion ensured prior to transport.

Antibiotics

Empirical treatment if infection is suspected

Patients with evidence of infection such as fever, leukocytosis, intravenous (IV) drug use, or a known infectious source should be started on empirical antibiotics after blood and urine cultures are drawn.

- Draw blood cultures and ESR
 - An ESR of less than 20 has excellent sensitivity for excluding a diagnosis of spinal epidural abscess
- Start empirical antibiotics with broad spectrum coverage
 - Abscesses are often multi-microbial
 - MRSA coverage – start vancomycin
 - Gram negative coverage – start third or fourth generation cephalosporin
 - Anaerobic coverage with metronidazole (should be considered in post-surgical and penetrating trauma)

Imaging Not Available

No MRI or CT

Without imaging, consider empirical medical treatment until imaging can be obtained.

- If the history, clinical picture suggests infection and epidural abscess is a possibility, start empirical antibiotics.
- If there is a history of cancer and neoplastic spinal cord compression is a possibility, start empirical corticosteroids.
- Expedited transfer to a facility with imaging capability is necessary.

Infectious Lesion

Empirical antibiotics and ESR

Suppurative infections of the spinal epidural space can cause neurological injury directly by compressing the spinal cord, or indirectly by compromising blood flow. The classic triad of fever, back pain, and neurologic dysfunction is not seen in most patients, and other symptoms may include localized back pain, radiculopathy, weakness, sensory changes, and sphincter dysfunction.

- STAT consultation with a spine surgeon or facilitation of transfer if none is available.
- Draw blood cultures and ESR
 - An ESR of less than 20 has excellent sensitivity for excluding a diagnosis of spinal epidural abscess
- Start empirical antibiotics with broad spectrum coverage
 - Abscesses are often multi-microbial
 - MRSA coverage – start vancomycin
 - Gram negative coverage – start third or fourth generation cephalosporin
 - Anaerobic coverage with metronidazole (should be considered in post-surgical and penetrating trauma)
- Antibiotics alone (without decompression) may be considered in patients who are either neurologically intact or who have had complete weakness for more than 48-72 hours.
 - In these patients, close observation for neurological worsening is advised given the high failure rates of medical management (6-49%)
 - Risk factors for neurological worsening
 - Diabetes
 - CRP > 115
 - WBC > 12 $10^3/\mu\text{L}$
 - age > 65
 - positive blood cultures
 - MRSA infection
- In patients with neurological deficits, early decompression, irrigation, and debridement is the mainstay of treatment [49].

MRI Spine

Spine imaging is available

Emergent MRI with gadolinium is the imaging modality of choice.

- CT with contrast and or CT myelogram is an alternative if MRI is contraindicated or not available.

Imaging is used to rule out any compressive etiology of the spinal cord like tumor, infection, or intervertebral disc herniation. It is important to communicate the neurological findings to your radiologist so that the proper location(s) of relevance is imaged.

- Quadriplegic patients should have at least the C-Spine imaged. Entire spine imaging (including the conus) may also be appropriate especially if the patient has known cancer.
- Paraplegic patients (if there are no symptoms in the arms) should have thoracic and lumbosacral spine imaged.
- A discussion with the radiologist is important to image the proper level, and to expedite the imaging so that treatments can be provided efficiently.
- It is important to notify the surgeon early when evaluating a patient with myelopathy, so that they are aware of timing of spine imaging and the potential need for surgical decompression.

Negative Imaging or Intrinsic Spinal Cord Lesion

ENLS Non-traumatic Weakness Protocol

Patients with acute symptoms of spinal cord dysfunction can have intrinsic or intra-medullary (non-compressive) abnormalities of the spinal cord on MRI. These include spinal infarct, inflammatory / demyelinating myelitis, infectious, and para-infectious disease.

Spinal cord ischemia may require further imaging with spinal angiography to identify arteriovenous malformation, arterial occlusion or other vascular abnormalities. Serum and CSF studies should be sent for evidence of autoimmune and infectious vasculitis. Serum studies should include serology for Lyme, syphilis, human immunodeficiency virus (HIV), and nutritional deficiencies including vitamin B12. CSF should be tested for viral studies including: herpes simplex virus 1,2 polymerase chain reaction (HSV PCR), cytomegalovirus (CMV) PCR, and Varicella Zoster Virus (VZV) IgM and IgG. Immunoglobulin indices should be evaluated including (but not limited to): IgG index, Albumin index, IgG synthesis rate and oligoclonal bands. Urine should be screened for copper deficiency (see the **ENLS Acute Non-Traumatic Weakness protocol**).

Pediatric considerations: In children with diffuse or progressive weakness, differential diagnosis should include metabolic disorders and genetic etiologies such as spinal muscular atrophy. For infants < 1 year of age, infant botulism should be considered, particularly if cranial nerves are involved or there is significant constipation. Treatment for this is with Botulism immune globulin (BabyBIG) and supportive care.

Negative spinal MRI in a patient with acute disturbance of motor and or sensory function (quadriplegia or paraplegia) should prompt evaluation for acute neuropathy, neuromuscular junction disorders and myopathy. Clinical presentation, CSF, and electrophysiologic studies may establish a diagnosis of Guillain-Barre syndrome (acute polyradiculoneuropathy), Myasthenia Gravis, Lambert-Eaton syndrome, or motor neuron disease (see **ENLS Acute Non-Traumatic Weakness protocol**).

Spinal cord injury without radiographic abnormality (SCIWORA) is a clinical diagnosis made in the setting of trauma and spinal cord dysfunction without an MRI abnormality. The cervical spinal cord is most likely affected due to increased mobility of the cervical spine. Children are high risk for this injury due to the relatively large head-to-body ratio in childhood. Treatment is non-surgical due to the absence of a surgical lesion. Supportive measures include spinal motion restriction with collars, braces, or orthosis for up to 3 months. The use of corticosteroids in this setting should be carefully evaluated with inherent risks and used judiciously.

Neoplastic Disease

Steroids, decompression, radiation oncology

Patients with malignant SCC and acute neurologic dysfunction should receive corticosteroids. Empirical treatment with corticosteroids is recommended when malignant SCC is suspected, even if unconfirmed by MRI spine imaging.

- Steroids are given to reduce edema and decrease the chance of spinal cord venous infarction. The use of steroids in patients with compression from epidural metastatic disease is considered part of standard medical therapy.
- Given the safety profile and efficacy of the lower dose of dexamethasone, we recommend a dexamethasone 10 mg IV loading dose followed by 4 mg oral/IV every 6 hours as maintenance.

Once imaging and diagnosis is confirmed, a combination of surgical treatment, radiotherapy, and chemotherapy is recommended. Surgical removal of tumor and spinal decompression is the primary and emergent treatment for malignant SCC.

- Early surgery (within 24 hours) with circumferential removal of the tumor, decompression of the spinal cord, and stabilization of the spine has been shown to significantly improve clinical outcomes (ambulation and pain), quality of life, and reduces need for narcotics and corticosteroids. It is also associated with a trend towards improved pain, quality of life and reduced need for analgesia.

Radiotherapy is used in conjunction with surgery and is recommended alone in patients with minimal neurologic dysfunction and high degree of tumor responsiveness to radiation. Hematologic tumors such as lymphoma, myeloma, and seminoma are highly radiosensitive, while breast, lung and prostate have intermediate radiosensitivity. There have been significant advancements in radiosurgical techniques, and all malignant SCC patients should be referred for evaluation.

Chemotherapy is not a mainstay for acute treatment of malignant SCC and is always delivered in conjunction with radiotherapy and surgery.

Paraplegia

Paraplegia is paralysis of the legs and lower body and is related to compression of the thoracic and/or lumbar spine.

Ventilatory issues are uncommon in patients with paraplegia/paraparesis, and transport and MRI imaging are safe.

Quadriplegia

Special airway issues

In the event of sudden or progressive quadriparesis or quadriplegia, the cause may be a cervical cord pathology. This may lead to hypoventilation because of both chest wall and diaphragmatic weakness, and respiratory assessment is vital before transport or MRI imaging.

Ventilatory issues are uncommon in patients with paraplegia/paraparesis, and transport and MRI imaging are safe.

Spinal Cord Compression

Suspected myelopathy

The presentation of spinal cord compression includes:

- Back/neck pain
- Bilateral weakness or paralysis of the limbs
- Urinary retention
- Obstipation
- Sensory level
 - Defined as a loss of sensation below the dermatomal level of compression.
- Spinal shock
 - Defined as flaccid paralysis, loss of reflexes, and sensation below the level of compression.
- Compression of the cauda equina can cause a similar clinical presentation with concomitant perineal (saddle) anesthesia and radicular pain.
- Spinal cord compression at or above T4 can lead to hemodynamic instability secondary to loss of sympathetic tone.
 - This leads to neurogenic shock with systemic hypotension and relative bradycardia.

SCC of the cervical segments can lead to quadriplegia, whereas compression below these levels causes paraplegia, which is far more common. Quadriplegia is the most disabling presentation of SCC, and when present in an alert and responsive patient, should suggest a cervical spinal cord lesion. The most widely used severity scale is the American Spinal Injury Association Impairment Scale (ASIA) which was devised for traumatic SCC.

Immediately after recognition or suspicion for SCC, spinal motion restriction should be instituted with a cervical collar and thoracolumbar motion restriction if warranted. Spinal motion restriction and precautions during patient care (and transport) prevent further cord compression and injury. While optimal blood pressure is not known, spinal cord blood flow is often impaired in the setting of a compressive lesion, and hypotension should be avoided.

Suspicion of Cancer

Possible metastasis

Consider spinal metastasis with spinal cord compression if there is a history of cancer, or new suspicion of cancer.

Suspicion of Infection

Possible epidural abscess

Suspicion for an infectious cause (epidural abscess) rises if the following are present:

- Fever
- Elevated WBC count
- Elevated ESR
- Diabetes
- History of intravenous drug use
- Known infectious source - current or past endocarditis, sepsis, chronic infection like osteomyelitis
- Any of the above with focal spine tenderness elicited by percussion (reflex hammer striking your finger placed over the vertebral spinous process)